

LA-UR-21-27522

Approved for public release; distribution is unlimited.

Exploring Gamma-Ray and Neutron Spectroscopy Missions to Mars Caves Title:

Berner, Angelica Marie Gasda, Patrick J. Author(s):

Mesick, Katherine Elizabeth

Hardgrove, Craig

Intended for: Student Symposium

Issued: 2021-07-30





Exploring Gamma-Ray and Neutron Spectroscopy Missions to Mars Caves

Angelica Berner (LANL, ASU)

Patrick Gasda (LANL), Katherine Mesick (LANL), Craig Hardgrove (ASU)

7/29/2021

Agenda

01


02

03

04

Background Project Results **Conclusions** Overview + Future Work Review of • Big-picture Description of Summary of look at the conclusions project simulation conception, project and setup and and next steps GRNS, and relevant configuration Description of Mars caves questions future work Overview of results





Background

- Mars Caves
 - Collapse chains → subsurface caves¹
 - Good candidates for habitability?
 - Potential to host water resources
 - Shielded from radiation




Image credit: Sauro, F., et al. (2020)Supplement

Sauro, F., et al. (2020). Lava tubes on Earth, Moon and Mars: A review on their size and morphology revealed by comparative planetology. Earth-Science Reviews. https://doi.org/10.1016/j.earscirev.2020.103288



Project Overview

 What: Investigating the sensitivity of active neutron detector systems in Mars caves

- How: Using MCNP6 to model neutron transport in different cave geometries, compositions, and detectors
- Why: To determine if active neutron measurements can be used to identify the hydration of cave walls vs floor vs ceilings, and to perform a concept study for potential future Mars cave missions in search of hydration and habitable regions




Results: Detectors on the Surface

- Box of air
 - Basic Mars atmosphere composition
- Box of planetary material
 - Crust+Mantle Mixture (Taylor, 2013)
- Detectors
 - Maintain same position from Curiosity model
 - Center of surface box




Results: Detectors on the Surface






Results: Simple Cave Model

- Geometry:
 - Hemisphere
- Composition:
 - Crust+Mantle mixture (Taylor, 2013)
 - Surrounding material is uniform
- Detectors:
 - Location varies (approaches wall)






Results: Simple Cave Model






Results: Ceiling H2O Content Varies (z=9)

 How does detector response change when only the top few meters of the cave ceiling change in composition?






Results: Ceiling H2O Content Varies (z=9)






Results: Ceiling H2O Content Varies (z=0)

 How does detector response change when the ceiling is defined as everything above z=0?






Results: Ceiling H2O Content Varies (z=0)






Results: Ceiling H2O Content Varies (z=0)






Results: Detectors on the Surface Near a Wall

- Wall added to surface model
 - 5m away, 6m tall





Results: Detectors on the Surface Near a Wall





Conclusions

- Initial results show that later time bins hold valuable information that allows us to discriminate between the ceiling and the wall of the cave
 - Particularly sensitive to composition and distance
- Immediate next steps:
 - Finish varying ceiling height runs
 - Explore and quantify the contribution of the wall to the cave response by varying detector distance on the surface
 - Look for possible second peak in real DAN data



Future Work

- Change the model geometry (steeper walls?)
- Change the detectors
- Acquire real data
 - BRAILLE team but with GRNS!

